

1. A planetary nebula is a

- a. shell of gases ejected from the surface of red giant star.
 - b. spherical, rapidly expanding cloud of gas produced by a supernova explosion.
 - c. cloud of gas surrounding a very young star in which planets are expected to form.
 - d. gas cloud surrounding a planet after its formation.
-

2. In what manner does an isolated white dwarf generate energy?

- a. An isolated white dwarf does not generate energy.
 - b. gravitational contraction
 - c. hydrogen fusion
 - d. helium fusion
-

3. What is a pulsar?

- a. Cepheid variable star with a period of a few days
 - b. pulsating white dwarf star, fluctuating rapidly in brightness
 - c. rapidly rotating neutron star, producing beams of radio energy and occasionally of X rays and visible light
 - d. very hot material orbiting a black hole
-

4. According to general relativity, why does Earth orbit the Sun?

- a. The Sun exerts a gravitational force on Earth across empty space.
 - b. Matter contains quarks, and Earth and the Sun attract each other with the “color force” between their quarks.
 - c. Space around the Sun is curved, and Earth follows a geodesic in this curved space.
 - d. Earth and the Sun are continually exchanging photons of light in a way that holds Earth in orbit.
-

5. What is the event horizon of a black hole?

- a. “surface” at which all events happen
 - b. “surface” at which any object passing through it will leave with greater energy than it entered
 - c. “surface” from inside of which nothing can escape
 - d. infinitesimally small volume at the center of the black hole that contains all of the black hole's mass
-

6. In the context of black holes, what is a wormhole?

- a. “tunnel” of undistorted space through an event horizon allowing objects to enter and leave a black hole without being torn apart
 - b. small, black hole through a solid object such as a planet
 - c. direct connection from any black hole to another part of spacetime
 - d. direct connection from a rotating black hole to another part of spacetime
-

7. Cepheid variable stars are useful to astronomers as indicators of

- a. white dwarf star behavior.
 - b. stars with very high-speed motion.
 - c. distance, particularly to nearby galaxies.
 - d. the existence of black holes.
-

8. Harlow Shapley first located the center of the Milky Way Galaxy in 1917 by

- a. measuring redshifts of stars in the galactic plane and disk.
 - b. observing the distribution of globular clusters in the galactic halo.
 - c. observing the distribution of hydrogen gas, measured by 21-cm radio emission.
 - d. measuring the positions of supernova explosions throughout the Galaxy.
-